arXiv:1012.0364v1 [quant-ph] 2 Dec 2010

Non-Markovian Quantum Trajectories of Many-Body Quantum Open Systems

Jun Jing[1][*], Xinyu Zhao[1], J. Q. You[2][†], Ting Yu[1][‡]
[1] *Center for Controlled Quantum Systems and Department of Physics and Engineering Physics, Stevens Institute of Technology, Hoboken, New Jersey 07030, USA*
[2] *Department of Physics and State Key Laboratory of Surface Physics, Fudan University, Shanghai 200433, China*

(Dated: December 3, 2010)

We have derived for the first time a set of non-Markovian quantum trajectory equations for many-body and multi-state quantum open systems. The exact time-local (time-convolutionless) quantum trajectory equations can be used to simulate quantum dynamics of coupled open systems interacting with a bosonic environment at zero temperature. Our general results are explained and illustrated with several atomic and optical open systems.

PACS numbers: 03.65.Ud, 03.67.Bg, 32.90+a

Introduction.—Dynamical aspect of quantum open systems has been investigated for many years and developed in different formalisms [1–9]. Typically, the state of an open system is described by a density operator governed by a master equation that plays a pivotal role in the fields of quantum optics and quantum information. When the environment is a structured medium or the system-environment interaction is strong, Lindblad Markov master equations are prone to fail badly. Then, non-Markovian master equations or their alternative non-Markovian quantum trajectory approaches must be used [10–14]. Despite extensive efforts, deriving a master equation without invoking the Born-Markov approximations (weak couplings and memoryless environment) has achieved only limited success in practice, and is mostly restricted in a few isolated models such as the quantum Brownian motion model [7], a leaky cavity model [15, 16], a phase damping model [17] and a two-level atom coupled to a quantized radiation field [18] etc. For a many-body open system such as an N-qubit system or a multi-state system coupled to a fully quantized environment, the existence of non-Markovian dynamic equations [19] such as master equations is still largely unknown.

It has been shown that the quantum state diffusion (QSD) equations for the pure states can formally solve a quantum open system coupled to a bosonic environment irrespective of environmental memory, coupling strength and the spectral density [20–24]. The generality of the QSD equation has given it appeal as both a numerical and theoretical tool for a non-Markovian open system. On the other hand, the time-nonlocal feature of the non-Markovian QSD equations has been a major obstacle in general implementations of the trajectory formalism for a realistic physical system. Therefore, it is highly desirable to develop a time-local non-Markovian formalism that is applicable to many-body open systems.

In this Letter, we report explicit constructions of exact time-local QSD equations for several many-body and multi-state quantum systems. Moreover, we show that these exact time-local QSD equations contain only polynomial noise terms. To the best of our knowledge, these time-local QSD equations for many-body open systems have never been derived before. Such non-Markovian quantum trajectory equations are capable of describing the transition from non-Markovian to Markov regimes for a genuine many-body quantum system.

The model and the quantum state diffusion equation.— A generic quantum open system in the system plus environment framework can be be written as (setting $\hbar = 1$):

$$H_{\rm tot} = H_{\rm sys} + H_{\rm int} + H_{\rm env} \qquad (1)$$

where $H_{\rm sys}$ is the Hamiltonian of the system of interest, $H_{\rm int} = \sum_{\mathbf{k}}(g_{\mathbf{k}}La^\dagger_{\mathbf{k}} + g^*_{\mathbf{k}}L^\dagger a_{\mathbf{k}})$ is the interaction Hamiltonian and $H_{\rm env} = \sum_{\mathbf{k}}\omega_{\mathbf{k}}a^\dagger_{\mathbf{k}}a_{\mathbf{k}}$ describes a quantized field (environment). Note that L is a system operator characterizing the mutual interaction between the system and the environment. At zero temperature $T = 0$, the environmental correlation function is determined by the noise operator $B(t) = \sum_{\mathbf{k}} g_{\mathbf{k}}a_{\mathbf{k}}e^{-\omega_{\mathbf{k}}t}$ in the interacting picture: $\alpha(t,s) = \langle 0|[B(t)+B^\dagger(t)][B(s)+B^\dagger(s)]|0\rangle = \sum_{\mathbf{k}}|g_{\mathbf{k}}|^2e^{-i\omega_{\mathbf{k}}(t-s)}$.

Here we use $|\Psi_{\rm tot}(t)\rangle$ to represent the state of the total system at time t, then the reduced density operator ρ_t for the system of interest is given by $\rho_t = {\rm Tr}_{\rm env}[|\Psi_{\rm tot}\rangle\langle\Psi_{\rm tot}|]$ obtained by tracing over the environmental degrees of freedom. If the system and its environment are initially uncorrelated, it has been shown that the density operator for the open system can be decomposed as a set of continuous quantum trajectories living in the system's Hilbert space, denoted by $\psi_t(z)$. The trajectory $\psi_t(z)$ is governed by a linear stochastic Schrödinger equation, termed linear QSD equation [20, 21]:

$$\frac{d}{dt}\psi_t(z) = \big(-iH_{\rm sys}+Lz_t-L^\dagger\int_0^t ds\alpha(t,s)\frac{\delta}{\delta z_s}\big)\psi_t(z) \qquad (2)$$

where $z_t = -i\sum_{\mathbf{k}} g_{\mathbf{k}}z^*_{\mathbf{k}}e^{i\omega_{\mathbf{k}}t}$ is a complex Gaussian process satisfying $M[z_t] = M[z_tz_s] = 0$, and $M[z^*_tz_s] =$

---

[*] Email address: Jun.Jing@stevens.edu

[†] Email address: jqyou@fudan.edu.cn

[‡] Email address: Ting.Yu@stevens.edu

$\alpha(t,s)$. Here $M[\cdot]$ denotes the ensemble average over the classical noise z_t. Note that $\alpha(t,s)$ is the correlation function defined above. The quantum trajectory $\psi_t(t)$ is designed to recover the density operator of the system by taking ensemble average: $\rho_t = M[|\psi_t(z)\rangle\langle\psi_t(z)|] = \int \frac{dz^2}{\pi} e^{-|z|^2} |\psi_t(z)\rangle\langle\psi_t(z)|$.

The Diósi-Gisin-Strunz QSD equation (2) is a remarkable result since it is completely general irrespective of spectral density of environment and coupling strength. The appearance of functional derivative in Eq. (2) is strongly reminiscent of the convolution kernel appearing in the Nakajima-Zwanzig master equations obtained from projection operator technique [25]. Thus, for the purpose of practical applications, it remains unresolved whether one can recast the existing functional derivative into a time-local form.

The ansatz to replace the functional derivative $\delta\psi_t(z)/\delta z_s = O(t,s,z)\psi_t(z)$ combining with the consistency condition $\frac{\delta}{\delta z_s}\frac{\partial \psi_t}{\partial t} = \frac{\partial}{\partial t}\frac{\delta \psi_t}{\delta z_s}$ yields the equation of motion of the O-operator [21–23]:

$$\frac{\partial O}{\partial t} = \left[-iH_{\rm sys} + Lz_t - L^\dagger \bar{O}, O\right] - L^\dagger \frac{\delta \bar{O}}{\delta z_s}. \qquad (3)$$

where $\bar{O}(t,z) \equiv \int_0^t \alpha(t,s)O(t,s,z)ds$ and the initial condition $O(s,s,z) = L$ is satisfied. Once the O-operator can be explicitly constructed, then the original non-local QSD equation (2) takes a time-local form:

$$\frac{d}{dt}\psi_t(z) = \left(-iH_{\rm sys} + Lz_t - L^\dagger \bar{O}(t,z)\right)\psi_t(z).$$

Clearly, finding the solutions to the nonlinear operator equation (3) is by no means trivial. Up to now, only a few isolated examples have been explicitly solved [21, 22, 26]. Here, we have derived the time-local QSD equations for a large class of many-body and multi-state systems that are of importance in quantum optics, atomic many-body physics and quantum information science [27].

For numerical simulations, one must use the normalized pure states $\tilde{\psi}_t(z) = \frac{\psi_t(z)}{||\psi_t(z)||}$ where $||\psi_t(z)|| = \langle\psi_t(z)|\psi_t(z)\rangle$, which is governed by a nonlinear version of the QSD equation:

$$\begin{aligned}\frac{d}{dt}\tilde{\psi}_t &= -iH_{\rm sys}\tilde{\psi}_t + \Delta_t(L)\tilde{z}_t\tilde{\psi}_t \\ &- \Delta_t(L^\dagger)\bar{O}(t,\tilde{z})\tilde{\psi}_t + \langle\Delta_t(L^\dagger)\bar{O}(t,\tilde{z})\rangle_t\tilde{\psi}_t,\end{aligned}$$

where $\Delta_t(A) \equiv A - \langle A\rangle_t$ for any operator A, $\langle A\rangle_t \equiv \langle\tilde{\psi}_t|A|\tilde{\psi}_t\rangle$ denotes the quantum average, and $\tilde{z}_t = z_t + \int_0^t \alpha^*(t,s)\langle L^\dagger\rangle_s ds$ is the shift noise. For all the numerical results to be presented below, we alway use the normalized nonlinear QSD equation.

Time-Local Non-Markovian QSD equations.— Equation (3) manifests that the O-operator may be determined by the commutation relations involving $H_{\rm sys}$ and L together with a set of basis operators.

In general, the O-operator may be expanded as [23],

$$O(t,s,z) = O^{(0)}(t,s) + \sum_{k=1}^{\infty} O^{(k)}(t,s,z) \qquad (4)$$

where $O^{(0)}(t,s) = \sum_j f_j(t,s)O_j^{(0)}$ is the noise-free term; $O^{(1)}(t,s,z) = \sum_j \int_0^t p_j^{(1)}(t,s,s_1)z_{s_1}ds_1 O_j^{(1)}$ is the linear-noise term, and in general $O^{(k)}(t,s,z) = \sum_j \int_0^t \cdots \int_0^t p_j^{(k)}(t,s,s_1,\cdots,s_k)z_{s_1}\cdots z_{s_k}ds_1...ds_k O_j^{(k)}$ contains kth-order polynomial noises. Note that all the basis operators $O_j^{(k)}$ and functions $f_j, p_j^{(k)}$ are noise-free. Moreover, the basis operators $O_j^{(k)}$ are time-independent. The equations for $f_j, p_j^{(k)}$ may be obtained from Eq. (3). Below, we shall show that the polynomial O-operators can be explicitly constructed for several many-body models ranging from discrete to continuous variable systems.

Theorem 1. (N-cavity model) *For the open system model (1) with* $H_{\rm sys} = \sum_{j=1}^N[\omega_j a_j^\dagger a_j + \lambda_j(a_j^\dagger a_{j+1} + a_j a_{j+1}^\dagger)]$ *(*$N+1=1$*) and* $L = \sum_{j=1}^N a_j$*, then the exact O-operator is noise-independent. Explicitly, it is given by* $O(t,s,z) = O^{(0)}(t,s) = \sum_{j=1}^N f_j(t,s)a_j$ *where* f_j *satisfy a set of differential equations [28].*

The proof and explicit construction of the O-operator for N-cavity model are straightforward (see [28]). Here, we provide an example of the O-operator with $N = 3$: $O(t,s,z) = f_1(t,s)a_1 + f_2(t,s)a_2 + f_3(t,s)a_3$, where $f_j(t,s)$, $(j = 1,2,3)$ satisfy the following equations: $\frac{\partial}{\partial t}f_j(t,s) = i\omega_j f_j(t,s) + \lambda_{j-1}f_{j-1}(t,s) + i\lambda_j f_{j+1}(t,s) + \sum_{k=1}^3 f_k(t,s)F_j(t)$ with the notation $F_j(t) = \int_0^t ds\alpha(t,s)f_j(t,s)$. The initial conditions are given by $f_1(s,s) = f_2(s,s) = f_3(s,s) = 1$.

Theorem 2. (Angular momentum model or N-level model) *For the open system model (1) with* $H_{\rm sys} = \omega J_z$ *and* $L = J_-$ *where* J_z *and* J_- *are angular momentum operators satisfying* $[J_z, J_\pm] = \pm J_\pm$, $[J_-, J_\pm] = \pm J_z$*, then the exact O-operator* $O(t,s,z)$ *can be explicitly constructed. More precisely, if* l *is the angular momentum quantum number (integer or half-integer), then the O-operator expressed in (4) has the basis operators* $O_j^{(k)} = J_z^{j-1}J_-^{k+1}$ *with* $k = 0,\cdots,2l-1$ *and* $j = 1,2,\cdots,2l-k$*.*

Let us emphasize that it is possible to construct the explicit O-operators for the multi-state systems with arbitrary energy spacings. The details of the proof for Theorem 2 can be found in [28]. It is instructive to provide a simple example of the O-operator of a spin-2 system (or a five-level system). Explicitly, the O-operator may be written as:

$$
\begin{aligned}
O(t,s,z) = & \sum_{j=1,2,3,4} f_j(t,s) O_j^{(0)} \\
& + \sum_{j=1,2,3} \int_0^t p_j^{(1)}(t,s,s_1) z_{s_1} ds_1 O_j^{(1)} \\
& + \sum_{j=1,2} \iint_0^t p_j^{(2)}(t,s,s_1,s_2) z_{s_1} z_{s_2} ds_1 ds_2 O_j^{(2)} \\
& + \iiint_0^t p_1^{(3)}(t,s,s_1,s_2,s_3) z_{s_1} z_{s_2} z_{s_3} ds_1 ds_2 ds_3 O_1^{(3)},
\end{aligned}
$$

where the time-independent basis operators $O_j^{(k)}$ are given by $O_j^{(0)} = J_z^{j-1} J_-$, $(j = 1, \cdots, 4)$; $O_j^{(1)} = J_z^{j-1} J_-^2, (j = 1, 2, 3)$, $O_j^{(2)} = J_z^{j-1} J_-^3, (j = 1, 2)$ and $O_1^{(3)} = J_-^4$. The differential equations governing the coefficients $f_j, p_j^{(k)}$ are explicitly given in [28].

Theorem 3. (N-qubit model) *For the open system model (1) with* $H_{\rm sys} = \frac{\omega}{2} \sum_j^N \sigma_z^{(j)}$ *and* $L = \sum_j^N \sigma_-^{(j)}$*, then the exact O-operator can be explicitly constructed, and contains* $(N-1)$*-order noises. That is,* $O_j^{(k)} = 0$ *if* $k \geqslant N$*.*

The general proof of Theorem 3 for an arbitrary N is rather cumbersome. Here we have proved Theorem 3 for $N \leqslant 7$ by explicit construction of the O-operators. If we use $m(N,k)$ $(0 \leqslant k \leqslant N-1)$ to denote the number of terms with k-fold noise integration in the O-operator, we have the following relations shown in Table I: $m(N,k) = m(N-1,k-1)$ and $m(N,0) = m(N-2,0) + N$ with $m(1,0) = 1$. Table I lists the numbers $m(k,N)$ for seven cases. Obviously, when $N = 1$, the O-operator just reduces to $O(t,s,z) = f(t,s)\sigma_-$ for a single qubit case [21, 23] with $H_{\rm sys} = \frac{\omega}{2}\sigma_z$ and $L = \sigma_-$. For $N = 2$, it is easy to check that the exact O-operator for the two coupled-qubit model has five terms: $O(t,s,z) = \sum_{j=1}^4 f_j(t,s) O_j^{(0)} + i \int_0^t p(t,s,s_1) z_{s_1} ds_1 O_1^{(1)}$. where $O_1^{(0)} = \sigma_-^A$, $O_2^{(0)} = \sigma_-^B$, $O_3^{(0)} = \sigma_z^A \sigma_-^B$, $O_4^{(0)} = \sigma_-^A \sigma_z^B$, and $O_1^{(1)} = \sigma_-^A \sigma_-^B$. As an application, we point out that the exact two-qubit time-local QSD equation allows us to calculate the entanglement evolution of the density matrix constructed from quantum trajectories [29]. Similarly, the exact three-qubit O-operator can be constructed explicitly (See [28]).

Our result in Theorem 3 regarding O-operators can readily be generalized to coupled N-qubit systems, which is expected to generate quantum coherence of qubit systems, including entanglement generation. The derivation is also applicable to the N-qubit systems where each qubit has a different frequency $H_{\rm sys} = \sum_j \omega_j \sigma_z^{(j)}/2$. Furthermore, N-qutrit or N-qudit models can be solved in a similar way.

N	$k=0$	$k=1$	$k=2$	$k=3$	$k=4$	$k=5$	$k=6$
1	1	0	0	0	0	0	0
2	2	1	0	0	0	0	0
3	4	2	1	0	0	0	0
4	6	4	2	1	0	0	0
5	9	6	4	2	1	0	0
6	12	9	6	4	2	1	0
7	16	12	9	6	4	2	1

TABLE I: This table summarizes the explicit constructions of the O-operator. The notation k stands for the orders of noises contained in the O-operator for an N-qubit dissipation model $(N = 1, 2, \cdots, 7)$. For $N = 2$, $m(2,0) = 2, m(2,1) = 1, m(2,k) = 0,\ (k \geqslant 2)$.

Numerical results and discussions.—Below, numerical results for the non-Markovian quantum dynamics of many-body systems are presented. For simplicity, and for recovering the Markov limit, we assumed that the correlation property of the noisy environment is described by the Ornstein-Uhlenbeck process: $\alpha(t,s) = \frac{\gamma}{2} e^{-\gamma|t-s|}$. Clearly, the Ornstein-Uhlenbeck noise approaches to the Markov limit when $\gamma \to \infty$. It should be noted that our exact time-local QSD equations are valid for arbitrary types of correlation functions.



FIG. 1: (Color online) Non-Markovian quantum interference of a four-level system (spin-3/2) with $H_{\rm sys} = \omega J_z$ and $L = J_-$. Initially, the system is in the state $\psi_0 = \frac{1}{2}(|1\rangle + |2\rangle + |3\rangle + |4\rangle)$, where $|j\rangle$'s, $j = 1, 2, 3, 4$ are the different levels for this four-level system. The time evolutions of coherences $|\rho_{23}|$ and $|\rho_{14}|$ and population terms ρ_{11} and ρ_{44} are calculated over 1000 trajectories. The blue solid curve stands for $\gamma = 0.3$, the red dashed curve for $\gamma = 1.0$ and the black dot-dashed curve for $\gamma = 3.0$.

The first example provides a simple picture of the effect of the non-Markovian atom-cavity interaction on quantum interference of a four-level atomic system. Both coherence decay and the population transition times from a higher level to a lower level are significantly different from the Markov limit as shown in Fig. 1. Our plots show that, in a non-Markovian regime, both coherence (Fig. 1a,1b) and population (Fig. 1c,1d) exhibit decay and revival patterns before eventually decaying to their final values. So the loss of coherence has effectively been suppressed due to the memory effect of the environment.



FIG. 2: (Color online) Evaluating fidelity under the influence of non-Markovian amplitude damping channel via quantum trajectories (1000 realizations). The initial state is the Werner state with parameter F: $W = \frac{F}{8} I_8 + (1-F)|\psi\rangle\langle\psi|$, where $|\psi\rangle = 1/\sqrt{3}(|100\rangle + |010\rangle + |001\rangle)$. (a) Fixed initial pure state with $F = 0$ for different memory time $\tau = 1/\gamma$; (b) Fidelity for different mixed states for the same channel with $\gamma = 0.3$.

The second example shows how a three-qubit system evolves when coupled to a common multiple-mode environment. The exact time-local QSD equation can be derived explicitly (For details, see [28]). Our numerical simulations with the zeroth-order $O^{(0)}(t,s)$ and the first-order terms $O^{(1)}(t,s,z)$ reveal the novel features of coherence dynamics measured by quantum fidelity. Fig. 3a shows the plot of fidelity against time for different environmental memory times $\tau = 1/\gamma, \gamma = 0.3, 1.0, 3.0$. Clearly, the shorter memory times give rise to better fidelity at time t. The result suggests a rather interesting feature that quantum coherence can typically survive longer in a non-Markovian dissipation environment. Another interesting feature arising from this four-level system is that, for a strong non-Markovian environment with $\gamma = 0.3$, the quantum fidelity is closely related to the degree of entanglement of initial three-qubit Werner states measured by the parameter F (Fig. 2b). As the initial state approaches separable state, the state becomes less affected by the environment reflecting the fragile feature of an entangled state [30].

Conclusion and outlook.—We have established the exact non-Markovian QSD equations for several important many-body and multi-state quantum systems including N-qubit systems and N-level atomic systems coupled to a common bosonic environment. The results can be extended to interacting qubits and high dimensional systems. Our findings will have many applications in many-body atomic coherence dynamics and quantum information science as illustrated by two examples in this Letter.

Our results can motivate other lines of research. Clearly, it is important to apply the exact many-body QSD equations to the important non-Markovian physical systems such as an atomic ensemble in an optical cavity and atomic dynamics in photonic crystals. Furthermore, it would be of interest to extend the current methods to the case of hybrid quantum systems consisting of both continuous and discrete variables.

Acknowledgments.—We thank J. H. Eberly and B. L. Hu for discussions. Grant support from DARPA QuEST No. HR0011-09-1-0008, the NSF No. PHY-0925174, the NBRPC No. 2009CB929300, and the NSFC Nos. 10625416, 10804069 is acknowledged.

[1] C. W. Gardiner and P. Zoller, *Quantum Noise* (Springer-Verlag, 2004).
[2] H. Carmichael, *An Open System Approach to Quantum Optics* (Springer, Berlin, 1994).
[3] J. Dalibard, Y. Castin, and K. Mølmer, Phys. Rev. Lett. **68**, 580 (1992).
[4] N. Gisin and I. C. Percival, J. Phys. A **25**, 5677 (1992); ibid. **26**, 2233 (1993).
[5] M. B. Plenio and P. L. Knight, Rev. Mod. Phys. **70**, 101 (1998).
[6] H. P. Breuer and F. Petruccione, *Theory of Open Quantum Systems* (Oxford, New York, 2002).
[7] B. L. Hu, J. P. Paz, and Y. Zhang, Phys. Rev. D **45**, 2843 (1992) and reference therein.
[8] A. Bassi, Phys. Rev. A **67**, 062101 (2003).
[9] L.-A. Wu, A. Bharioke, and P. Brumer, J. Chem. Phys. **129**, 041105 (2008).
[10] N. Vats and S. John, Phys. Rev. A **58**, 4168 (1998).
[11] H. P. Breuer, D. Faller, B. Kappler, and F. Petruccione, Phys. Rev. A **60**, 3188 (1999).
[12] P. Lambropoulos *et al.*, Rep. Prog. Phys **63**, 455 (2000).
[13] P. Borri *et al.*, Phys. Rev. Lett. **87**, 157401 (2001).
[14] V. Lopez-Richard *et al.*, Nano Lett. **9**, 3129 (2009).
[15] T. Yu, Phys. Rev. A **69**, 062107 (2004).
[16] H. N. Xiong, W. M. Zhang, X. G. Wang and M. H. Wu, Phys. Rev. A **82**, 012105 (2010).
[17] For example, see, M. Orszag, *Quantum Optics* (Springer, 2000).
[18] C. Anastopoulos and B. L. Hu, Phys. Rev. A **62**, 033821 (2002).
[19] For an open system in a weakly non-Markovian regime, a systematic post-Marov master equation can be derived, see T. Yu, L. Diósi, N. Gisin, and W. T. Strunz, Phys. Lett. A, **265**, 331 (2000).
[20] L. Diósi and W. T. Strunz, Phys. Lett. A **235**, 569 (1997).
[21] L. Diósi, N. Gisin and W. T. Strunz, Phys. Rev. A **58**, 1699 (1998).
[22] W. T. Strunz, L. Diósi and N. Gisin, Phys. Rev. Lett. **82**, 1801 (1999).
[23] T. Yu, L. Diósi, N. Gisin and W. T. Strunz, Phys. Rev. A **60**, 91 (1999).
[24] J. Gambetta and H. M. Wiseman, Phys. Rev. A **66**, 012108 (2002).
[25] F. Haake, *Quantum Statistics in Optics and Solid State Physics* (Springer-Verlag, Berlin, 1973).
[26] J. Jing and T. Yu, Phys. Rev. Lett. to be published (2010).
[27] M. A. Nielson and I. L. Chuang, *Quantum Computa-*

tion and Quantum Information (Cambridge Univ. Press, 2000).

[28] See supplementary material.

[29] X. Zhao *et al.* (to be submitted).

[30] T. Yu and J. H. Eberly, Quantum Inf. Comput. **7**, 459 (2007).